

Here, living with ovarian cancer. Here for the good, the bad, and the quiet spaces in between. Here for you. Here for me. Here.

Your guide to treatment with **ELAHERE**

In a clinical study, median overall survival was 16.5 months for ELAHERE vs 12.7 months for traditional chemotherapy.

What is ELAHERE?

ELAHERE is a prescription medicine used to treat adults with folate receptor-alpha positive ovarian cancer, fallopian tube cancer, or primary peritoneal cancer who:

- have not responded to or are no longer responding to treatment with platinum-based chemotherapy and
- have received 1 to 3 prior types of chemotherapy.

Your healthcare provider will perform a test to make sure that ELAHERE is right for you.

It is not known if ELAHERE is safe and effective in children.





Patient portrayal.

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about ELAHERE?

ELAHERE can cause serious side effects, including:

Eye problems. Eye problems are common with ELAHERE and can also be severe. Tell your healthcare provider right away if you develop any eye problems during treatment with ELAHERE, including blurred vision, dry eyes, sensitivity to light, eye pain, eye redness, or new or worsening vision changes. **Continue reading information about eye problems on page 2.**



H TABLE OF CONTENTS

This guide is designed to help you understand what to expect during treatment with ELAHERE, from getting started to support along the way.

About ELAHERE	3
Results	6
Possible side effects	8
Caring for your eyes	10
Starting ELAHERE	12
Talking to your doctor	14
ELAHERE Support Services	15
Key terms	17



Not intended for pediatric patients. Patient portrayal.

IMPORTANT SAFETY INFORMATION (CONT'D)

- Your healthcare provider will send you to see an eye care professional to check your eyes
 before you start treatment with ELAHERE, during treatment with ELAHERE, and as needed
 for any worsening signs and symptoms of eye problems.
- Your healthcare provider will prescribe steroid eye drops and lubricating eye drops before
 you start and during your treatment with ELAHERE. You should use eye drops as directed
 by your healthcare provider.
- Do not wear contact lenses throughout your treatment with ELAHERE unless you are told to use them by your healthcare provider.



ABOUT ELAHERE

Is ELAHERE right for me?

ELAHERE is the **first targeted therapy** developed and approved for **FRa+ platinum-resistant ovarian cancer**.

Ask your doctor about ELAHERE to see if it may be right for you.



You have received 1 to 3 prior types of chemotherapy

ELAHERE is for people who have already tried chemotherapy and may need a change in their treatment approach.



Your ovarian cancer is platinum resistant

Platinum-resistant ovarian cancer is when your cancer comes back or progresses after receiving a type of chemotherapy that contains platinum.



Your cancer is positive for a type of protein called FRa

If your cancer is FRa+, you may be eligible for targeted treatment with ELAHERE.

Key terms: <u>ovarian cancer</u>; <u>FRα+</u>.

FRa=folate receptor alpha.

IMPORTANT SAFETY INFORMATION (CONT'D)

What should I tell my healthcare provider before receiving ELAHERE?

Tell your healthcare provider about all of your medical conditions, including if you:

- have vision or eye problems.
- have numbness or tingling in your hands or feet.
- have liver problems.
- are pregnant or plan to become pregnant. ELAHERE can harm your unborn baby.
 Tell your healthcare provider right away if you become pregnant or think you may be pregnant during treatment with ELAHERE.



ABOUT ELAHERE

Explore a targeted treatment option



What is FRa and why is it important?

Folate receptor alpha, or FRa, is a protein often found in people with ovarian cancer. Your doctor may refer to it as a biomarker. Testing for FRa can help your doctor determine if ELAHERE is right for you.

How is ELAHERE designed to target and kill cancer cells?

ELAHERE may affect cancer cells as well as certain healthy cells in the body, which can cause side effects.

ELAHERE is a type of treatment called an antibody-drug conjugate (ADC). ADCs are different from traditional chemotherapies. They are designed to attack cancer in a targeted way.



ELAHERE attaches to the outside of the cell



After binding to FRa, ELAHERE is taken into the cell



ELAHERE releases a strong cancerkilling drug inside the cell



The cancer-killing drug can also impact other nearby cells that may or may not be cancerous


Key term: biomarkers.

IMPORTANT SAFETY INFORMATION (CONT'D)

Patients who are able to become pregnant:

- Your healthcare provider should do a pregnancy test before you start treatment with FLAHERE.
- You should use an effective birth control (contraception) during treatment and for 7 months after your last dose of ELAHERE.
- are breastfeeding or plan to breastfeed. It is not known if ELAHERE passes into your breast milk. Do not breastfeed during treatment and for 1 month after your last dose of ELAHERE.





In a clinical study, median overall survival was 16.5 months for ELAHERE vs 12.7 months for traditional chemotherapy.



Patient portrayal.

IMPORTANT SAFETY INFORMATION (CONT'D)

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Taking certain other medicines during treatment with ELAHERE may cause side effects.



RESULTS

ELAHERE may help stop cancer from progressing

In a clinical study of 453 people with FRa+ platinum–resistant ovarian cancer, ELAHERE helped extend the time to disease progression compared with traditional chemotherapy.

Median progression-free survival



More time without the cancer growing or spreading with ELAHERE

Median overall survival

Some people taking ELAHERE lived longer

In the same clinical study, ELAHERE was proven to help people live longer compared with traditional chemotherapy.

minetuninal surentansine-gym injection 100 mg (227 people)

MONTHS

Traditional chemotherapy (226 people)

Key terms: median progression-free survival; median overall survival; traditional chemotherapy.

IMPORTANT SAFETY INFORMATION (CONT'D)

What are the possible side effects of ELAHERE?

ELAHERE can cause serious side effects, including:


• **Eye problems.** Eye problems are common with ELAHERE and can also be severe. Tell your healthcare provider right away if you develop any eye problems during treatment with ELAHERE, including blurred vision, dry eyes, sensitivity to light, eye pain, eye redness, or new or worsening vision changes.



RESULTS

Some people had a better response with ELAHERE

The study also found that more people's tumors shrank by at least 30% or disappeared when treated with ELAHERE vs when treated with traditional chemotherapy. The decrease in tumor size lasted a median of 6.8 months in those treated with ELAHERE and 4.5 months in those treated with traditional chemotherapy.



^{*}In the clinical study, 225 people treated with ELAHERE and 224 people treated with traditional chemotherapy were eligible to be evaluated for overall response.

Key term: overall response.

IMPORTANT SAFETY INFORMATION (CONT'D)

Lung problems (pneumonitis). ELAHERE can cause severe or life-threatening
inflammation of the lungs that may lead to death. Tell your healthcare provider right
away if you get new or worsening symptoms, including trouble breathing, shortness
of breath, cough, or chest pain.



POSSIBLE SIDE EFFECTS

What are the possible side effects of treatment with ELAHERE?

ELAHERE can cause serious side effects, including:



Eye problems

Eye problems are common with ELAHERE and can also be severe.

Tell your healthcare provider right away if you develop any eye problems during treatment with ELAHERE, including blurred vision, dry eyes, sensitivity to light, eye pain, eye redness, or new or worsening vision changes.



Lung problems (pneumonitis)

ELAHERE can cause severe or life-threatening inflammation of the lungs that may lead to death.

Tell your healthcare provider right away if you get new or worsening symptoms, including trouble breathing, shortness of breath, cough, or chest pain.



Peripheral neuropathy

Nerve problems called peripheral neuropathy are common with ELAHERE and can also be severe.

Your healthcare provider will monitor you for signs and symptoms of nerve problems. Tell your healthcare provider if you get new or worsening numbness, tingling, burning sensation or pain in your hands or feet or muscle weakness.



POSSIBLE SIDE EFFECTS

What are the possible side effects of treatment with ELAHERE? (cont'd)



The most common side effects and abnormal lab results include:

- Increased liver enzymes in the blood
- Feeling tired
- Blurred vision
- Nausea
- Diarrhea
- Stomach-area (abdominal) pain
- Changes in the cornea (part of the eye)
- Peripheral neuropathy
- Muscle, bone, or joint pain

- Decreased red or white blood cell counts
- Decreased platelets
- Decreased magnesium level in the blood
- Dry eye
- Constipation
- Vomiting
- Decreased albumin level in the blood
- Decreased appetite

Your healthcare provider may change your dose of ELAHERE, delay treatment, or completely stop treatment if you have certain side effects.

Due to side effects in a clinical study, 24% of people reported serious adverse events, 9% of people stopped treatment, 54% of people had dose delays, and 34% of people had their dose reduced.

Of those treated with ELAHERE in a clinical study, 1% experienced hair thinning or hair loss.



These are not all of the possible side effects of ELAHERE. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088.



CARING FOR YOUR EYES

How could ELAHERE affect my eyes?

People treated with ELAHERE can develop eye-related side effects, which are common and can be severe. These may include blurred vision, dry eyes, sensitivity to light, eye pain, eye redness, new or worsening vision changes, temporary changes to the surface of the eye, or cataract. Tell your doctor if you have a history of vision or eye problems, or if you experience any of the above eye problems during treatment.



Your doctor will have you meet with an eye doctor before your first infusion and periodically throughout your treatment.



Eye-related side effects most commonly occurred between the **second**and third infusion

of ELAHERE.



1% of people in the study stopped treatment because of eye problems.



Eye care during treatment

Visit <u>elahere.com/eye-care</u> to learn about managing eye-related side effects.

IMPORTANT SAFETY INFORMATION (CONT'D)

Peripheral neuropathy. Nerve problems called peripheral neuropathy are common
during treatment with ELAHERE and can also be severe. Your healthcare provider will
monitor you for signs and symptoms of nerve problems. Tell your healthcare provider if
you get new or worsening numbness, tingling, burning sensation or pain in your hands
or feet or muscle weakness.




CARING FOR YOUR EYES

How can I care for my eyes with eye drops?

Your doctor will prescribe **steroid** eye drops and over-the-counter, **lubricating**, preservative-free eye drops for use as directed before and during treatment with ELAHERE. Unless your doctor tells you otherwise, avoid contact lenses throughout treatment.

Steroid eye drops



Lubricating eye drops



The use of preservative-free lubricating eye drops is also recommended at least 4 times daily and as needed during treatment. Wait at least 10 minutes after applying steroid eye drops before using lubricating eye drops.

IMPORTANT SAFETY INFORMATION (CONT'D)

The most common side effects and abnormal labs of ELAHERE include:

- increased liver enzymes in the blood
- feeling tired
- blurred vision
- nausea
- diarrhea
- stomach-area (abdominal) pain
- changes in the cornea (part of the eye)
- peripheral neuropathy
- muscle, bone, or joint pain

- · decreased red or white blood cell counts
- decreased platelets
- decreased magnesium level in the blood
- dry eye
- constipation
- vomiting
- decreased albumin level in the blood
- · decreased appetite



STARTING ELAHERE

Steps to help you start treatment

Before you start treatment

FRa

Your doctor will test a tissue sample to confirm your cancer is FRa+ and help determine if ELAHERE may be right for you.

To do this, they can use tumor tissue stored from an earlier biopsy or surgery or take a new sample.



Your doctor will also have you meet with an eye doctor before your first infusion and periodically throughout your treatment.

This may help you manage any eye-related side effects.



Your doctor will prescribe steroid eye drops and advise you on the use of lubricating eye drops before and during treatment with ELAHERE.

You should use steroid and lubricating drops as directed by your doctor.



Need reminders to use your eye drops?

There are several mobile apps available to help you and/or your caregiver remember your eye drop schedule. Check them out at your preferred app store.

Before each dose of ELAHERE, you will receive medicines to help prevent infusion-related reactions, nausea, and vomiting.

- Some of these medicines may be taken orally and some may be given to you by infusion into your vein
- Usually, these medicines will be given on the same day as the infusion. However, your doctor may choose
 to give them to you the day before you start ELAHERE

Tell your doctor about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Taking ELAHERE with certain other medicines may cause side effects.

IMPORTANT SAFETY INFORMATION (CONT'D)

Your healthcare provider may change your dose of ELAHERE, delay treatment, or completely stop treatment if you have certain side effects.

These are not all of the possible side effects of ELAHERE. Call your doctor for medical advice about side effects. You are encouraged to report side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1–800–FDA–1088.



STARTING ELAHERE

How will I receive ELAHERE?

ELAHERE is given by infusion (IV or port) every 3 weeks



ELAHERE dosage is determined by your weight.

To ensure accurate dosing, your doctor will weigh you prior to treatment. If you have side effects with ELAHERE, your doctor may reduce your dose, delay your dose, or completely stop treatment.

IV=intravenous.

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about ELAHERE?

ELAHERE can cause serious side effects, including:

Eye problems. Eye problems are common with ELAHERE and can also be severe. Tell your healthcare provider right away if you develop any eye problems during treatment with ELAHERE, including blurred vision, dry eyes, sensitivity to light, eye pain, eye redness, or new or worsening vision changes.

- Your healthcare provider will send you to see an eye care professional to check your eyes before you start treatment with ELAHERE, during treatment with ELAHERE, and as needed for any worsening signs and symptoms of eye problems.
- Your healthcare provider will prescribe steroid eye drops and lubricating eye drops before
 you start and during your treatment with ELAHERE. You should use eye drops as directed
 by your healthcare provider.
- Do not wear contact lenses throughout your treatment with ELAHERE unless you are told to use them by your healthcare provider.



TALKING TO YOUR DOCTOR

Here to start a conversation about ELAHERE

It's important to ask questions and start a conversation with your doctor about what is next in your treatment journey. The list of questions below can help guide your conversation.

Not sure where to start the conversation with your doctor? Try asking:

What is FRa, and why is it important?
Have I been tested for FRa?
What were my results, and how do they affect my treatment path?
What is platinum resistance?
If platinum-based chemotherapy stops working for me, what are my next steps?
How does ELAHERE work differently from other treatments?
What is the goal of treatment with ELAHERE?
Can you talk me through what to expect if I start treatment with ELAHERE?



Build your Discussion Guide

Visit <u>elahere.com/talking-to-your-doctor</u> to personalize a discussion guide that can help you start a conversation with your doctor.

IMPORTANT SAFETY INFORMATION (CONT'D)

What should I tell my healthcare provider before receiving ELAHERE?

Tell your healthcare provider about all of your medical conditions, including if you:

- have vision or eye problems.
- have numbness or tingling in your hands or feet.
- have liver problems.



ELAHERE SUPPORT SERVICES

ELAHERE Support Services (ESS) is here to help you navigate your treatment journey

Once enrolled, ESS can help you with insurance coverage questions and provide nursing support for your ELAHERE questions. Simply talk with your doctor to submit the ELAHERE Support Services enrollment form.

Nurse Navigators offered through ELAHERE Support Services

Trained nurses are available by phone to answer questions about FDA-approved information on ELAHERE while offering ongoing support throughout your treatment journey.

Case Managers offered through ELAHERE Support Services

Case Managers will help answer your questions about insurance coverage, copay, or the Patient Assistance Program (PAP).

Financial assistance includes the following options:



If you have commercial insurance*:

Patients eligible for the ESS Co-pay Assistance Program could pay as little as \$0 for their medication.



If you have no insurance or your current insurance does not provide sufficient coverage or treatment:

The Patient Assistance Program provides eligible patients with ELAHERE at no charge.[†]

IMPORTANT SAFETY INFORMATION (CONT'D)

Tell your healthcare provider about all of your medical conditions, including if you:

 are pregnant or plan to become pregnant. ELAHERE can harm your unborn baby. Tell your healthcare provider right away if you become pregnant or think you may be pregnant during treatment with ELAHERE.





IMPORTANT SAFETY INFORMATION (CONT'D)

Patients who are able to become pregnant:

- Your healthcare provider should do a pregnancy test before you start treatment with ELAHERE.
- You should use an effective birth control (contraception) during treatment and for 7 months after your last dose of ELAHERE.
- are breastfeeding or plan to breastfeed. It is not known if ELAHERE passes into your breast milk. Do not breastfeed during treatment and for 1 month after your last dose of ELAHERE.



KEY TERMS

Key terms to keep in mind as you learn about ELAHERE

Biomarkers are genes, proteins, or other substances that can be tested, found, and measured to give your doctor more information about your cancer—like whether the tumor is growing or how well you may respond to certain treatments. (page 4)

FRa+ (folate receptor alpha positive) is a biomarker test result that means you have a certain amount of FRa on the surface of your cancer cells. (page 3)

Median overall survival is the length of time that half of the people were alive after starting treatment. (page 6)

Median progression-free survival is the length of time that the treatment stops the cancer from growing or getting worse for half of the people. A median is the middle number in a group of numbers that are listed from lowest to highest. (page 6)

Ovarian cancer is used throughout this guide to refer to ovarian cancer, fallopian tube cancer, and primary peritoneal cancer. (page 3)

Overall response is a measure of all people who had either a complete or partial response. (page 7)

Traditional chemotherapy in the clinical study included anticancer drugs that are not targeted and do not contain a material called platinum. (page 6)

IMPORTANT SAFETY INFORMATION (CONT'D)

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Taking certain other medicines during treatment with ELAHERE may cause side effects.



For all the reasons there are to be here, ask your doctor about **ELAHERE**

In a clinical study, median overall survival was 16.5 months for ELAHERE vs 12.7 months for traditional chemotherapy.

Visit elahere.com to learn more.

Need support or have questions?

Call 1-833-ELAHERE (1-833-352-4373)

Monday to Friday, 8:00 AM to 8:00 PM ET

or email ELAHERESupport@cardinalhealth.com

*Eligibility: Available to patients with commercial insurance coverage for **ELAHERE** who meet eligibility criteria. This co-pay assistance program is not available to patients receiving prescription reimbursement under any federal, state, or government-funded insurance programs (for example, Medicare [including Part D], Medicare Advantage, Medigap, Medicaid, TRICARE, Department of Defense, or Veterans Affairs programs) or where prohibited by law. Offer subject to change or termination without notice. Restrictions, including monthly maximums, may apply. This is not health insurance. For full Terms and Conditions, visit elahere.com/copay-terms or call 1-833-352-4373 for additional information. To learn about AbbVie's privacy practices and your privacy choices, visit https://abbv.ie/corpprivacy.

[†]Criteria include: patients who are uninsured or have insurance that excludes coverage for ELAHERE (including patients on Medicare or Medicaid), residents of the United States or Puerto Rico, and patients who meet the financial eligibility requirements. Terms and conditions apply.

IMPORTANT SAFETY INFORMATION (CONT'D)

What are the possible side effects of ELAHERE?

ELAHERE can cause serious side effects, including:

Eye problems. Eye problems are common with ELAHERE and can also be severe. Tell your
healthcare provider right away if you develop any eye problems during treatment with
ELAHERE, including blurred vision, dry eyes, sensitivity to light, eye pain, eye redness, or
new or worsening vision changes.

